

Catalog
HL - UDCqPM2X

Description
1.3ml

Store at: 20°C
PI/HL UDCqPM2X-01


Product description

Dcore qPCR Mix with Uracil DNA Glycosylase (UDG) is a premixed 2X concentrated reagent ready to use for Real-time Polymerase chain reactions (PCR). The premix 2x reagent inclusive of HotStart Taq Polymerase Enzyme, UDG enzyme, and dNTPs helps in an easy experimental setup and avoids nonspecific amplifications. Template of choice along with primer probe mix is to be added such that the final concentration of the PCR Mix is 1X in the final assay. The Dcore qPCR mix with UDG is designed for TaqMan-based assays with DNA templates for contamination-free amplifications.

The reagent along with HotStart Taq Polymerase Enzyme, dNTPs come with dUTP and Uracil DNA Glycosylase (UDG) enzyme. The incorporation of dUTP in place of dTTP to the growing strand during PCR makes it an ideal degradation target for the UDG enzyme. The UDG enzyme gets inactivated at higher temperatures and therefore does not interfere with the amplification while clearing out any potential contamination that may be carried over from previous experiments. The mix is available with and without UDG.

Features

- Contains HotStart Taq Polymerase Enzyme to provide a better yield with 5'-3' Polymerase and Exonuclease activity. Contains UDG enzyme to prevent carry-over contaminations
- Includes HotStart Taq Polymerase reaction buffer, dNTPs for ready to use PCR applications
- High specificity and sensitivity
- Buffer enhancements guarantee performance and reliability



Reagents Supplied

Dcore qPCR Mix (2X) with UDG

Storage conditions

Dcore qPCR Mix (2X) with UDG should be stored at -20°C

Recommended reaction set-up for PCR

The Dcore PCR mix works the best for specific primers and is not recommended for random primers!

Prepare PCRs using required volumes of freeze-thawed components in the PCR hood as recommended in the table below

PCR Protocol

Component	Volume(µL)
Dcore qPCR Mix with UDG	13.0
Forward primer (10 pm/µL)	0.4-0.7
Reverse primer (10 pm/µL)	0.4-0.7
TaqMan Probe	Variable (User defined)
Template DNA or cDNA	Variable (User defined)
Nuclease Free Water	Variable (User defined)
Total Volume	26

The reaction set up is for guidance and it can be modified according to the user's need.

Reference Guide

The current product is sufficient for 100 reactions if the above protocol is used.

Note: Mix the vials with the reaction set up by gentle tapping. A short spin is recommended after gentle mixing to ensure that the reagent mix is not sticking to the walls of the PCR tube

Thermocycling Protocol

Place the tubes in the Thermal Cycler and start the polymerase chain reaction protocol. Below is a general template for PCRs and should be optimized for good results

Recommended PCR Program

Operation	Temp	Time	Cycles
Initial denaturation	95 °C	15 min	1
Denaturation	95 °C	15 sec	35-40 cycles
Annealing	60 °C	60 sec	

Applications

- Routine PCR reactions
- Gene expression analysis
- Amplified fragment length polymorphism
- cDNA amplification
- Molecular identification of organisms
- Viral load detection
- Pharmacogenomics
- Single nucleotide polymorphisms

Other PCR products from Huwel LifeSciences that you may be interested

S.No.	Product description	Catalogue No.
1.	Dcore qPCR Mix (2X) ***	HL-DCqPM2X
2.	Taq Polymerase Enzyme (5U/μl)	HL - Taq - 50 - 250Units HL - Taq - 100 - 500Units HL - Taq - 200 - 1000Units
3.	HotStart Taq Polymerase Enzyme (5U/μl)	HL - HSTaq - 50 - 250Units HL - HSTaq - 100 - 500Units HL - HSTaq - 200 - 1000Units
4.	QuickStart Taq Polymerase Enzyme (5U/μl)	HL - QSTaq - 50 - 250Units HL - QSTaq - 100 - 500Units HL - QSTaq - 200 - 1000Units

*** This PCR mix is without UDG

For further information on protocols and details, please contact our technical support:
info@huwelllifesciences.in



HLSS Manufacturing Pvt Ltd,
 Plot Nos; M 14, M 15, M 16,
 TSIIC, Medical Devices Park,
 Sultanpur Village, Ameenpur Mandal,
 Sangareddy Dist, Telangana-502319,
www.huwelllifesciences.in, Email:info@huwelllifesciences.in